

| SEDICO WELL | | | | | | 1. | |
|-------------|-------------|---|----------------------|----------------------|---------|----|--|
| Chronost | Lithostrat. | Forams and Nannos bioevents | Foraminifer Biozones | Nannofossil Biozones | Remarks | | |
| | | Top well | | | | | |
| | | 100m | | | | | |
| | | 200m | | | | | |
| | | 300m | | | | | |
| | | 320m ← <i>M. aragonensis</i> (266) | E9 | NP 15 | ? | ? | Not sampled |
| | | 400m ↑ <i>G. mexicana kugleri</i> and <i>Nannotetrina</i> sp. (383) | | | | | The two bioevents are recorded in the same sample probably due to the low sampling frequency. |
| | | 500m | | | | | |
| | | 600m | | | | | |
| | | 600m ↑ <i>D. sublodoensis</i> (591) | E8 | NP 14 | | | This correlation doesn't agree with the standard biostratigraphic schemes probably because the recorded events are displaced by caving. |
| | | 600m ↓ <i>A. soldadoensis</i> (591) | | | | | Other event: HO of <i>Acarinina soldadoensis</i> (591). |
| | | 600m ↑ <i>G. higginsi</i> (639) | | | | | |
| | | 700m | | | | | |
| | | 800m | | | | | |
| | | 900m ↑ <i>D. lodoensis</i> (883) | E6-E7 | NP 12-NP 13 | | | The foraminifer E7/E8 Zonal boundary is uncertain because <i>G. nuttalli</i> is absent and <i>G. higginsi</i> occurs very rarely only in samples 639 and 591. |
| | | 900m ↓ <i>M. subbotinae</i> (883) | | | | | The nannofossil NP 12-NP 13 Zones are indistinct because the HO of <i>Tribrachiatus orthostylus</i> , probably for poor preservation, is recorded below the LO of <i>Discoaster lodoensis</i> . |
| | | 990m ↑ <i>M. aragonensis</i> (994) | E5 | NP 11 | | | The foraminifer E6-E7 Zones are indistinct because <i>Acarinina cuneicamerata</i> is absent. |
| | | 990m ↑ <i>T. orthostylus</i> (1008) ↑ <i>M. formosa</i> (1008) | | | | | Other event: HO of <i>Morozovella formosa</i> (639). |
| | | 990m ↓ <i>F. tympaniformis</i> (1032) | | | | | |
| | | 1000m ↑ <i>M. velascoensis</i> (1066) | E4 | | | | This correlation doesn't agree with the standard biostratigraphic schemes probably because the recorded events are displaced by caving. |
| | | 1000m ↓ | E3 | NP 10 | | | Other bioevent: HO (highest occurrence) of <i>Fasciculithus tympaniformis</i> (1032). |
| | | 1050m ↑ <i>D. multiradiatus</i> (1150) | E1-E2 | NP 9 | | | The foraminifer P5-E2 Zones are indistinct because <i>Acarinina sibaiyaensis</i> is absent and <i>P. wilcoxensis</i> first appears above in sample 1008, probably for poor preservation. |
| | | 1050m ↓ <i>G. pseudomenardii</i> (1150) | P 5 | | | | Other bioevents: LOs of <i>Morozovella subbotinae</i> (1150), <i>Sphenolithus anharopus</i> (1172) and <i>Acarinina soldadoensis</i> (1208). The latter marks the base of the foraminifer P4c Subzone. |
| | | 1050m ↑ <i>H. kleinpelli</i> (1208) | P 4 | NP 6-NP 8 | | | Other bioevent: LO (lowest occurrence) of <i>Igorina albeari</i> (1228). |
| | | 1050m ↑ <i>G. pseudomenardii</i> and <i>F. tympaniformis</i> (1216) | | NP 5 | | | This horizon subdivides the foraminifer P 3 Zone into two Subzones (a and b). |
| | | 1200m ← <i>M. angulata</i> and <i>C. pelagicus</i> (1248) | P 3 | NP 4 | | | |
| | | 1210m bottom depth | | | | | |
| PALEOCENE | Late | Vena d'Oro Marl | | | | | |
| | | Scaglia Rossa | | | | | |
| | | 1200m | | | | | |
| | | 1210m | | | | | |
| | | 1250m bottom depth | | | | | |