

L. LIMONTA, P. DIOLI, N. BONOMELLI

Heteroptera on flowering spontaneous herbs in differently managed orchards (*)

Abstract - Heteroptera were collected on spontaneous herbs, during flowering, in two environments with different mowing management. The highest number of Heteroptera was collected on *Medicago sativa*, *Daucus carota*, *Centaurea nigrescens* and *Erigeron annuus*. As foreseeable, a higher number of species was found in the environment less influenced by traditional farming methods and with a higher number of plant species.

Riassunto - Eterotteri su piante spontanee fiorite in colture con diversa gestione.

Sono state considerate le specie di Eterotteri raccolte su piante spontanee, al momento della fioritura, in due ambienti con diversa gestione degli sfalci. Le piante su cui sono stati catturati più Eterotteri sono *Medicago sativa*, *Daucus carota*, *Centaurea nigrescens* ed *Erigeron annuus*. Come prevedibile, la località caratterizzata da un maggiore numero di specie è risultata quella con una maggiore ricchezza botanica, meno disturbata dalle pratiche culturali.

Key words: Heteroptera, herbs, mowing management.

As Heteroptera show a considerable differentiation in food diets, they can offer a good measure of biodiversity; the richness in species in this Order is strongly related to the total insect diversity (Duelli & Obrist, 1998). In many crops Heteroptera represent most of the total fauna; they are particularly abundant in unsprayed apple and pear orchards, where they form 30-50% of predators throughout the growing season (Fauvel, 1999). Many species are supported by moss and lichens, found on unsprayed trees harbouring bark dwelling insects, such as Psocids (Fauvel, 1999).

Fencerows and hedgerows, called by Altieri (1999) "*biological corridors*", provide overwintering sites, food and alternative preys for parasitoids and predators, increasing abundance and favouring the diffusion in the crops (Remund *et al.*, 1992).

(*) Work published with a grant of Regione Lombardia and E.C. Research Project: "Exploitation of olive growing in Lombardy (n. 528/99).

Heteroptera present in two different plant mixtures were considered in a previous work (Limonta *et al.*, 2003). In this work Heteroptera collected on flowering spontaneous herbs in two differently managed crops were studied, as different mowing management can influence the composition of Heteroptera fauna (Di Giulio *et al.*, 2001). In fact, a careful management is important; it was observed that mowing all the plot in some perennial plants, like *Medicago sativa* L., causes the destruction of the field fauna while strip harvesting creates a more suitable environment and favours moving of beneficial insects to crop plants (van den Bosch *et al.* 1967; Summers, 1976; van den Bosch *et al.*, 1982).

MATERIALS AND METHODS

Researches were carried out in olive groves in Manerba del Garda and Portese (Brescia province) near Garda Lake.

In the first place unsprayed olive groves with ground cover plants, in "Parco Naturale della Rocca", were surveyed. Herbs comes from previous seeding associated to olive grove: wide areas of meadow Fabaceae like clover, lucerne, five-finger, and also of Poaceae are present; however planted seeds are mingled with spontaneous ones, that belong to Apiaceae, Geraniaceae, Ranunculaceae and Asteraceae.

In Portese the olive grove is clean cultivated, only in the edge there are plants, regularly mowed. The ground is harrowed on the row and on the inter-row. It borders on another olive grove with ground cover regularly mowed. Both are pruned and sprayed with copper fungicide and organophosphates.

Insect sampling

Insects were sampled weekly, from April to September, from 1999 to 2001, on herbs under or near olive trees; the samples were collected on flowering plants, covering at least a square meter surface. In table 1 Families and species of collected plant are listed. In Parco della Rocca in Manerba a high number of plant species is present, but only synanthropic, ruderal or cultivated plants covered an adequate surface to sample.

Samples were collected with a suction device (a garden Blower-vac, whose direction of rotation was reversed). Sampling on plants lasted one minute each.

The exclusion of plants with a reduced flowering surface precluded the use of biodiversity indexes.

RESULTS AND DISCUSSION

In table 2 the species of Heteroptera and the number of collected individuals are listed; diet, pabulum and corology are shown in table 3 (Dioli, 1980, 1997; Tamanini, 1988; Southwood & Leston, 1959). The highest biodiversity is observed in Manerba olive grove, where the prevailing species, *Orius niger*, predator, form little more than 20% of

Table 1 - Plants surveyed in Portese and in Manerba.

Family	Species	Portese	Manerba
Caryophyllaceae	<i>Silene vulgaris</i> (Moench) Garke	*	
Fabaceae	<i>Lotus corniculatus</i> L.		*
	<i>Medicago sativa</i> L.	*	*
	<i>Trifolium repens</i> L.		*
Cistaceae	<i>Helianthemum nummularium</i> (L.) spp. <i>obscurum</i> (Celak) Ho		*
Apiaceae	<i>Daucus carota</i> L.	*	*
Rubiaceae	<i>Galium lucidum</i> All.		*
	<i>Galium verum</i> L.		*
Convolvulaceae	<i>Convolvulus arvensis</i> L.	*	
Verbenaceae	<i>Verbena officinalis</i> L.	*	
Lamiaceae	<i>Ajuga reptans</i> L.		*
	<i>Centaurea nigrescens</i> Willd.	*	*
	<i>Calamintha selvetica</i> Bromf.	*	*
	<i>Mentha</i> sp.	*	*
Scrophulariaceae	<i>Veronica chamaedrys</i> L.		*
Asteraceae	<i>Bellis perennis</i> L.	*	*
	<i>Erigeron annuus</i> (L.) Pers.	*	*
	<i>Taraxacum officinale</i> Web.		*

population, while in Portese *Nysius graminicola*, phytophagous, make up 80% and the total number of species is less than half of the number in the other place.

Species of Heteroptera and host plants, on which insects were collected, are reported in tables 4 and 5. In Portese *Erigeron annuus*, *Medicago sativa* and *Centaurea nigrescens* were the most visited plants; in Manerba *M. sativa* was colonized by 20 species of Heteroptera, *Daucus carota* and *C. nigrescens* by 11 species. *E. annuus* was studied by many Authors as nectar source, with contrasting results: *Pimpla turionellae* L. is unable to exploit nectar (Wackers *et al.*, 1996), while the opposite happens with *Ooencyrtus nezarae* Ishii (Takeshi & Numata, 2000); Fleischer *et al.* (1988) demonstrated that the presence of the plant decreased the number of adults of *Lygus lineolaris* (Palisot de Beauvois) in cotton fields. *Daucus carota* has exposed and accessible nectars (Patt *et al.*, 1997), exploited by many parasitoids (Idris & Grafius, 1997; Manojlovic *et al.*, 2000, 2001); *M. sativa* has a high nectar-carbohydrate production (Davis, 2001) and enhance the level of parasitoids (Lawrence *et al.*, 2003). *C. nigrescens* is present in uncultivated ground and permanent meadows; it is considered a weed (Ward *et al.*, 1974; Solymosi, 1996) and it is neglected as potential nectar source because it has hidden nectaries, like all Asteraceae, on which parasitoids can forage only difficultly (Patt *et al.*, op. cit.).

Table 2 - Species of Heteroptera and number of specimen collected in the two places.

		Portese	Manerba
Anthoridae	<i>Orius niger</i> Wolff	169	195
Nabidae	<i>Aptus mirmicoides</i> (O.G.Costa)		5
	<i>Nabis pseudoferus</i> Remane	2	16
	<i>Nabis punctatus</i> A.Costa	8	28
Miridae	§ <i>Deraeocoris punctulatus</i> (Fallén) *		13
	<i>Deraeocoris serenus</i> Douglas & Scott	16	16
	<i>Halticus apterus</i> (L.)*	4	19
	<i>Adelphocoris lineolatus</i> (Goeze)	25	169
	<i>Adelphocoris seticornis</i> (Fabricius)		12
	<i>Charagochilus gyllenhalii</i> (Fallén)	1	57
	<i>Liocoris tripustulatus</i> (Fabricius)*		1
	<i>Orthops kalmi</i> (L.)		3
	<i>Stenodema laevigatum</i> (L.)		1
	<i>Stenotus binotatus</i> (Fabricius)	8	2
	<i>Trigonotylus ruficornis</i> (Goeffroy)	13	5
	<i>Chlamydatus pullus</i> Reuter	3	2
	<i>Criocoris crassicornis</i> (Hahn)		79
	<i>Plagiognathus arbustorum</i> (F.)	11	2
Tingidae	<i>Copium clavicorne</i> (L.)		1
	<i>Kalama tricornis</i> (Schränk)	1	
	<i>Tingis crespata</i> (Herrich-Schäffer)		1
	<i>Tingis auriculata</i> (A.Costa)		3
Alydidae	<i>Alydus calcaratus</i> (L.)		7
	<i>Camptopus lateralis</i> (Germar)		4
Coreidae	<i>Syromastes rhombeus</i> (L.)		3
	<i>Coriomeris denticulatus</i> (Scopoli)	1	2
Rhopalidae	<i>Rhopalus subrufus</i> (Gmelin)	5	6
	<i>Stictopleurus crassicornis</i> (L.)		2
Lygaeidae	<i>Heterogaster affinis</i> Herrich-Schäffer	1	
	<i>Nysius graminicola</i> (Kolenati)	285	50
	<i>Nysius senecionis</i> (Schilling)		1
	<i>Macroplox preyssleri</i> (Fieber)		1
	<i>Oxycarenus pallens</i> (Herrich-Schafer)	1	3
	<i>Beosus quadripunctatus</i> (O.F. Müller)	3	14
	<i>Megalonotus sabulicola</i> (Thomson)		3
	<i>Paromius gracilis</i> (Rambur)		1
	<i>Peritrechus gracilicornis</i> Puton		1
	<i>Rhyparochromus vulgaris</i> (Schilling)		1
	<i>Scolopostethus thomsoni</i> Reuter		4
Cydnidae	<i>Legnotus picipes</i> (Fallén)	2	
Pentatomidae	<i>Dolycoris baccarum</i> (L.)		4
	<i>Eysacoris aeneus</i> (Scopoli)		1
	<i>Nezara viridula</i> (L.)		21
	<i>Piezodorus lituratus</i> (Fabricius)		1
	<i>Graphosoma lineatum</i> (L.)		7
	<i>Vilpianus galii</i> (Wolff)		2

* species collected on mixed flowering plants Diet, pabulum and chorology of species are reported in tab. 3; species in normal character were already listed in a previous paper (Limonta *et al.*, 2003).

§ *Deraeocoris punctulatus* is separated from *D. serenus*, light colored, even if Wagner (1975) considered the last one the only valid species.

Table 3 - Main characteristics of Heteroptera surveyed in Portese and in Manerba.

Classification	Diet	Pabulum	Chorology ¹	Portese	Manerba
Anthocoridae					
<i>Orius niger</i> Wolff, 1811	Zoophagous	Prey Insects and Mites on <i>Artemisia</i> , <i>Achillea</i> , <i>Verbascum</i>	Mediterranean	*	*
Nabidae					
<i>Aptus mirmicoides</i> (O.G. Costa, 1834)	Zoophagous	Prey little Insects	Centraleuropean		*
<i>Nabis pseudoferus</i> (Remane, 1949)	Zoophagous	Prey Insect adults and larvae	N-Anatolian-European	*	*
<i>Nabis punctatus</i> (A. Costa, 1847)	Zoophagous	Prey Insects	European-Saharian	*	*
Miridae					
<i>Deraeocoris punctulatus</i> (Fallén, 1807)	Zoophytophagous	Prey little Insects and eggs	Siberian-European		*
<i>Deraeocoris serenus</i> Douglas & Scott, 1868	Zoophytophagous	Prey little Insects	Mediterranean	*	*
<i>Halticus apterus</i> (L., 1758)	Phytophagous	<i>Ononis</i> , <i>Galium</i>	Mediterranean	*	*
<i>Adelphocoris lineolatus</i> (Goeze, 1778)	Phytophagous	<i>Artemisia</i> , <i>Verbascum</i> , <i>Achillea</i>	European	*	*
<i>Adelphocoris seticornis</i> (Fabricius, 1775)	Phytophagous	Fabaceae: <i>Trifolium</i> , <i>Vicia</i>	Asiatic-European		*
<i>Charagochilus gyllenhalii</i> (Fallén, 1807)	Phytophagous	<i>Galium</i> spp.	Palearctic	*	*
<i>Liocoris tripustulatus</i> (Fabricius, 1781)	Phytophagous	<i>Parietaria</i> spp., <i>Solanum</i> spp., <i>Urtica</i> , <i>Verbascum</i> spp., <i>Salvia</i>	Siberian-European		*
<i>Orthops kalmi</i> (L., 1758)	Phytophagous	Apiaceae: <i>Daucus</i> and <i>Pastinaca</i>	Palearctic		*
<i>Stenodema laevigatum</i> (Linnaeus, 1758)	Phytophagous	Poaceae	Holarctic		*
<i>Stenotus binotatus</i> (Fabricius, 1794)	Phytophagous	Asteraceae, mainly <i>Dactylis</i> , <i>Phleum pratense</i>	Holarctic	*	*
<i>Trigonotylus ruficornis</i> (Geoffroy, 1758)	Phytophagous	Poaceae	Holarctic	*	*
<i>Chlamydatus pullus</i> Reuter, 1870	Phytophagous	<i>Achillea</i> , <i>Trifolium</i>	Palearctic	*	*
<i>Criocoris crassicornis</i> (Hahn, 1834)	Phytophagous	<i>Galium</i> sp.	Turanian-European-Mediterranean		*
<i>Plagiognathus arbustorum</i> (Fabricius, 1794)	Phytophagous	<i>Urtica</i> sp.	Palearctic	*	*

¹ According to Vigna Taglianti *et al.* (1999).

(Table 3 continued)

Classification	Diet	Pabulum	Chorology	Portese	Manerba
Tingidae					
<i>Copium clavicorne</i> (Linneus, 1758)	Phytophagous	<i>Teucrium chamaedris</i>	S-European		*
<i>Kalama tricornis</i> (Schränk)	Phytophagous	<i>Artemisia campestris</i>	Siberian-European	*	
<i>Tingis crespata</i> (Herrich-Schäffer, 1838)	Phytophagous	<i>Hieracium</i> , <i>Artemisia</i>	Centraleuropean Mediterranean		*
<i>Tingis auriculata</i> (A. Costa, 1847)	Phytophagous	<i>Daucus</i> spp., <i>Caucalis platycarpus</i> , <i>Cnidium silaifolium</i> , <i>Torilis</i> spp.	S-European		*
Alydidae					
<i>Alydus calcaratus</i> (L., 1758)	Phytophagous	Different plants	Holarctic		*
<i>Camptopus lateralis</i> (Germar, 1817)	Phytophagous	Polyphagous	S-Palearctic		*
Coreidae					
<i>Syromastes rhombeus</i> (Linneaus, 1767)	Phytophagous	<i>Euphorbia</i> , Caryophyllaceae	Centraleuropean		*
<i>Coriomeris denticulatus</i> (Scopoli, 1763)	Phytophagous	Polyphagous	Siberian-European Mediterranean	*	*
Rhopalidae					
<i>Rhopalus subrufus</i> (Gmelin, 1790)	Phytophagous	<i>Trifolium</i> , <i>Urtica</i> , <i>Salvia pratensis</i>	Cosmopolitan	*	*
<i>Stictopleurus crassicornis</i> (L., 1758)	Phytophagous	<i>Abutilon</i> , <i>Artemisia</i> , <i>Achillea</i>	European		*
Lygaeidae					
<i>Heterogaster affinis</i> Herrich-Schäffer, 1835	Detritivorous	<i>Cytisus scoparius</i> , <i>Artemisia</i> sp., <i>Coronilla</i> sp., <i>Scabiosa</i> sp.	Turanian-European Mediterranean	*	
<i>Nysius graminicola</i> (Kolenati, 1845)	Detritivorous Phytophagous	Different plants	Mediterranean	*	*
<i>Nysius senecionis</i> (Schilling, 1829)	Detritivorous Phytophagous	<i>Artemisia</i>	European Mediterranean		*
<i>Macroplox preyssleri</i> (Fieber, 1837)	Phytophagous	<i>Thymus serpyllum</i>	European		*
<i>Oxycarenus pallens</i> (Herrich-Schäffer, 1850)	Phytophagous	<i>Centaurea</i>	Centralasiatic Mediterranean	*	*

(Table 3 continued)

Classification	Diet	Pabulum	Chorology	Portese	Manerba
<i>Beosus quadripunctatus</i> (O.F. Müller, 1766)	Detritivorous	<i>Cytisus scoparius</i> , <i>Silene</i> sp., <i>Erica arborea</i> , <i>Calluna vulgaris</i>	Turanian Mediterranean	*	*
<i>Megalonotus sabulicola</i> (Thomson, 1870)	Detritivorous	Asteraceae, Poaceae	European		*
<i>Paromius gracilis</i> (Rambur, 1839)	Phytophagous	<i>Elanthus</i> , <i>Imperata</i> , <i>Andropogon</i>	Mediterranean-Ethiopian		*
<i>Peritrechus gracilicornis</i> Puton, 1877	Detritivorous	Poaceae, Ericaceae	Turanian-Mediterranean		*
<i>Rhyparochromus vulgaris</i> (Shilling, 1829)	Detritivorous	Poaceae	Turanian-European		*
<i>Scolopostethus thomsoni</i> Reuter, 1874	Detritivorous Phytophagous	<i>Urtica</i>	Holarctic		*
Cydnidae					
<i>Legnotus picipes</i> (Fallén, 1807)	Phytophagous	<i>Artemisia</i> spp	Siberian-European	*	
Pentatomidae					*
<i>Dolycoris baccarum</i> (L., 1758)	Phytophagous	Polyphagous ¹	Asiatic-European		*
<i>Eysacoris aeneus</i> (Scopoli, 1763)	Phytophagous	Lamiaceae	Asiatic-European	*	
<i>Nezara viridula</i> (Linnaeus, 1758)	Phytophagous	Polyphagous	Afrotropical-Indoaustral Mediterranean		*
<i>Piezodorus lituratus</i> (Fabricius, 1794)	Phytophagous	Fabaceae	Turanian-European Mediterranean		*
<i>Graphosoma lineatum</i> (Linnaeus, 1758)	Phytophagous	Apiaceae	Paelearctic		*
<i>Vilpianus galii</i> (Wolff, 1802)	Phytophagous	<i>Galium verum</i>	Turanian- S-European Mediterranean		*

Table 4 - Plants and Heteroptera collected in Portese olive grove.

Heteroptera \ Plants		<i>Silene vulgaris</i> (Moench) Gark	<i>Medicago sativa</i> L.	<i>Daucus carota</i> L.	<i>Convolvulus arvensis</i> L.	<i>Verbena officinalis</i> L.	<i>Calamintha sylvatica</i> Bromf.	<i>Mentha</i>	<i>Erigeron annuus</i> (L.) Pers.	<i>Bellis perennis</i>	<i>Centaurea nigrescens</i> Willd.
Anthocoridae	<i>Orius niger</i> Wolff		x	x	x		x	x	x		x
Nabidae	<i>Nabis pseudoferus</i> Remane								x		
	<i>Nabis punctatus</i> A.Costa		x						x		
Miridae	<i>Deraeocoris serenus</i> (Douglas & Scott)		x				x		x		
	<i>Halticus apterus</i> (L.)								x		x
	<i>Adelphocoris lineolatus</i> (Goeze)	x	x		x				x		x
	<i>Charagochilus gyllenhalii</i> (Fallén)*		x								x
	<i>Stenotus binotatus</i> (Fabricius)										x
	<i>Trigonotylus ruficornis</i> (Goeffroy)	x	x			x					x
	<i>Chlamidatus pullus</i> Reuter	x	x								
	<i>Plagiognathus arbustorum</i> (Fabricius)	x									x
Tingidae	<i>Kalama tricornis</i> (Schrank)				x						
Coreidae	<i>Coriomeris denticulatus</i> (Scopoli)								x		
Rhopalidae	<i>Rhopalus subrufus</i> (Gmelin)					x	x	x	x		
Lygaeidae	<i>Heterogaster affinis</i> Herrich-Schäffer							x			
	<i>Nysius graminicola</i> (Kolenati)	x	x	x	x				x	x	
	<i>Oxycarenus pallens</i> (Herrich-Schäffer)			x							
	<i>Beosus quadripunctatus</i> (O.F.Müller)							x			
Cydnidae	<i>Legnotus picipes</i> (Fallén)			x							x
total		5	8	4	4	2	3	4	9	1	8

* the species was collected on mixed flowering plants.

In both places (tabb. 4-5), the minute pirate bug *Orius niger*, predator of insects and mites, was frequently collected on different plants. The presence in both environments is due to the ability of Anthocoridae to tolerate pesticides, probably thanks to the high mobility and to the large number of generations (Fauvel, 1999). Other predators present in both places are damsel bugs *Nabis pseudoferus*, species not influenced by intensive management of grass (Di Giulio *et al.*, 2001), and *N. punctatus*, lit-

Table 5 - Plants and Heteroptera collected in Manerba olive grove.

Heteroptera \ Plants		<i>Medicago sativa</i> L.	<i>Trifolium repens</i> L.	<i>Lotus corniculatus</i> L.	<i>Helianthemum nummularium</i> (L.) Miller ssp. <i>obscurum</i>	<i>Daucus carota</i> L.	<i>Galium verum</i> L.	<i>Galium lucidum</i> All.	<i>Ajuga reptans</i> L.	<i>Calamintha selvalica</i> Bromf.	<i>Mentha</i>	<i>Veronica chamaedrys</i> L.	<i>Erigeron annuus</i> (L.) Pers	<i>Bellis perennis</i> L.	<i>Centaurea nigrescens</i> Willd.	<i>Taraxacum officinale</i> Web.
Anthracoridae	<i>Orius niger</i> Wolff	x	x		x					x	x				x	
Nabidae	<i>Aptus mirmicoides</i> (O.G.Costa)				x		x	x								
	<i>Nabis pseudoferus</i> Remane	x	x							x						
	<i>Nabis punctatus</i> A.Costa	x	x			x	x					x			x	
Miridae	<i>Deraeocoris serenus</i> Douglas & Scott	x	x			x	x						x			
	<i>Deraeocoris punctulatus</i> (Fallén)*															
	<i>Halticus apterus</i> (L.)*	x				x									x	
	<i>Adelphocoris lineolatus</i> (Goeze)	x	x			x	x	x			x				x	
	<i>Adelphocoris seticornis</i> (F.)	x													x	
	<i>Charagochilus gyllenhalii</i> (Fallén)	x	x				x						x			x
	<i>Liocoris tripustulatus</i> (F.)*	x	x					x								
	<i>Orthops kalmi</i> (L.)					x									x	
	<i>Stenodema laevigatum</i> (L.)		x													
	<i>Stenotus binotatus</i> (Fabricius)		x		x			x								
	<i>Trigonotylus ruficornis</i> (Goeffroy)	x	x			x										
	<i>Chlamydatus pullus</i> Reuter							x								
	<i>Criocoris crassicornis</i> (Hahn)						x									
	<i>Plagiognathus arbustorum</i> (F.)								x							
Tingidae	<i>Copium clavicornis</i> (L.)				x											
	<i>Tingis crespata</i> (Herrich-Schäffer)								x							
	<i>Tingis auriculata</i> (A.Costa)		x	x												
Alydidae	<i>Alydus calcaratus</i> (L.)	x											x			
	<i>Camptopus lateralis</i> (Germar)	x														
Coreidae	<i>Syromastus rhombeus</i> (L.)					x						x				
	<i>Coriomeris denticulatus</i> (Scopoli)	x														
Rhopalidae	<i>Rhopalus subrufus</i> (Gmelin)	x					x								x	
	<i>Stictopleurus crassicornis</i> (L.)															
Lygaeidae	<i>Nysius graminicola</i> (Kolenati)	x				x				x				x		
	<i>Nysius senecionis</i> (Schilling)															x
	<i>Macroplox preyssleri</i> (Fieber)							x								
	<i>Oxycarenus pallens</i> (Herrich-Schafer)					x									x	
	<i>Beosus quadripunctatus</i> (O.F.Müller)	x													x	
	<i>Megalonotus sabulicola</i> (Thomson)														x	
	<i>Paromius gracilis</i> (Rambur)				x											
	<i>Peritrechus gracilicornis</i> Puton								x							
	<i>Rhyparochromus vulgaris</i> (Schilling)							x								
	<i>Scolopostethus thomsoni</i> Reuter	x														
Pentatomidae	<i>Dolycoris baccarum</i> (L.)	x						x							x	
	<i>Eysacoris aeneus</i> (Scopoli)															
	<i>Nezara viridula</i> (L.)	x								x						
	<i>Piezodorus lituratus</i> (Fabricius)	x														
	<i>Graphosoma lineatum</i> (L.)					x										
	<i>Vilpianus galii</i> (Wolff)						x	x								
total		20	3	9	4	11	8	7	5	4	2	2	3	1	11	2

* the species was collected on mixed flowering plants.

the specialized predator (Chiappini & Reguzzi, 1994); both are spread all over Italy on herbs in dry environments. Also the plant bug *Deraeocoris serenus* is widely spread in Italy. *Aptus mirmicoides* and *Deraeocoris punctulatus* are the species present only in Manerba. The damsel bug *A. mirmicoides*, collected on different plants, more common in woods, preys little insects; Maceljski & Balarin (1977) list it among predators of *Corythuca ciliata* (Say), emphasising the lesser efficiency with low temperatures, while Buchholz & Schruft (1994) report it as predator of eggs, larvae and pupae of *Eupoecilia ambiguella* Hbn.. *D. punctulatus* preys little insects on different plants and trees; it was collected on mixed flowering of *Galium verum* and *Silene vulgaris*.

Mirids are more abundant in Manerba and in fact they seem to be more linked to the host plant and more susceptible to the effect of chemical sprays than anthocorids (Fauvel, 1999).

Among phytophagous mirids, *Halticus apterus* is a widely common species, that colonize *Medicago sativa*, other Fabaceae and *Galium*; it is worth mentioning the collection of individuals on *C. nigrescens* and *E. annuus*. It was collected also on *D. carota* mixed with *M. sativa*. It is not influenced by intensive management of grassland like *Adelphocoris lineolatus*, *Stenodema laevigatum* and *Dolycoris baccarum* (Di Giulio *et al.*, 2001). Few individuals of the last two species were found only in Manerba (tab. 2).

Charagochilus gyllenhalii, palearctic species widespread in Italy, is linked to *Galium*, but it was collected also on Fabaceae and Asteraceae.

The specimen of *Liocoris tripustulatus*, with the same diffusion of the previous species, was collected in Manerba on mixed flowering of *Lotus*, *Medicago* and *Galium*; usually it colonize *Parietaria*, *Solanum*, *Urtica*, *Verbascum* and *Salvia*.

Also *Peritrechus gracilicornis* was found only in Manerba; in fact the species is negatively affected by intensive management, like *Adelphocoris seticornis* (Di Giulio *et al.*, 2001).

Adelphocoris lineolatus, *A. seticornis*, *Stenotus binotatus*, *Trigonotylus ruficornis*, *Nysius graminicola*, *Nezara viridula*, phytophagous, are common species and they can be pests of various crops (Southwood & Leston, 1959; Balarin, 1978; Di Giulio *et al.*, 2001; Accinelli *et al.*, 2002; Every *et al.*, 1992; Bin & Colazza, 1986; Parenzan *et al.*, 1994; Tanaka *et al.*, 1991). Even *Criocoris crassicornis* and *Beosus quadripunctatus* were frequently collected, but they are harmless: the first one colonize *Galium*, the second one is detritivorous.

On the whole, the higher plant biodiversity and the scarce mowing in Manerba directly affect the Heteropterofauna that is more abundant in term of number of species and of individuals than in Portese, where grass management is intensive.

REFERENCES

- ACCINELLI G., FERRARI R., DRADI D., POZZATI M., 2002 - Caratteristiche distintive, danni e difesa: i principali miridi fitofagi di interesse agrario. - *Informatore agrario*, 58 (41): 47-50.
- ALTIERI M.A. 1999 - The ecological role of biodiversity in agroecosystems. - *Agriculture, Ecosystems and Environment* 74: 19-31.

- BALARIN I., 1978 - Prilog poznavanju Heteroptera u fauni lucerista jadranskog podruca. - Poljoprivredna Znanstvena Smotra, 45 (55): 119-129 (english summary).
- BIN F. COLAZZA S., 1986 - Esplosione demografica di *Nysius graminicola* Kol. sul sorgo in Italia centrale (Rincoto Ligeide). - Informatorere agrario, 42 (34): 75-77.
- BUCHHOLZ U., SCHRUFF G., 1994 - Räuberische Arthropoden auf Blüten und Früchten der Weinrebe (*Vitis vinifera* L.) als Antagonisten des Einbindigen Traubenwicklers (*Eupoecilia ambiguella* Hbn.) (Lep., Cochylidae). - J. Appl. Ent., 118: 31-37.
- CHIAPPINI E., REGUZZI M.C., 1994 - Osservazioni su *Nabis punctatus* Costa (Rhynchota Nabidae). - Redia, LXXVII (1): 79-99.
- DAVIS A.R., 2001 - Searching and breeding for structural features of flowers correlated with high nectar-carbohydrate production. - Acta Horticulturae, 561: 107-121.
- DI GIULIO M., EDWARDS P.J., MEISTER E., 2001 - Enhancing insect diversity in agricultural grasslands: the roles of management and landscape structure. - J. Appl. Ecol., 38: 310-319.
- DIOLI P., 1980 - Eterotteri della brughiera di Rovasenda (Piemonte). - Estratto dai Quaderni sulla "Struttura delle zoocenosi terrestri". 1. La brughiera pedemontana III. Roma, Collana del P.F. CNR "Promozione della qualità dell'ambiente": 35-56.
- DIOLI P., 1997 - Gli Eterotteri del Monte Barro. Studi geobotanici ed entomofaunistici del Parco Regionale del Monte Barro. - Memorie della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano. 27 (2): 159-174.
- DUELLI P., OBRIST M.K., 1998 - In search of the best correlates for local organismal biodiversity in cultivated areas. - Biodiversity and Conservation, 7: 297-309.
- EVERY D., FARREL J.A., STUFKENS M.W., 1992 - Bug damage in New Zealand wheat grain: the role of various heteropterous insects. - N. Z. Jl crop hort. Sci., 20 (3): 305-312.
- FAUVEL G., 1999 - Diversity of Heteroptera in agroecosystems: role of sustainability and bioindication. - Agriculture, Ecosystem and Environment, 74: 275-303.
- FLEISCHER S.J., GAYLOR M.J., HUE N.E., 1988 - Dispersal of *Lygus lineolaris* (Heteroptera: Miridae) adults through cotton following nursery host destruction. - Env. Ent., 17 (3): 533-541.
- FORMANN R.T.T., GORDON M., 1986 - Landscape ecology. - Ed. Wiley & Sons, New York, pp 620.
- IDRIS A.B., GRAFIUS E., 1997 - Nectar-collecting behaviour of *Diadegma insulare* (Hymenoptera: Ichneumonidae), a parasitoid of diamondback moth (Lepidoptera: Plutellidae). - Env. Ent., 26 (1): 114-120.
- KHAMRAEV A. SH., 2001 - Miridae bugs on cotton. Zashchita i Karantin Rastenii, 10: 44-45.
- LAWRENCE L., SCHELLHORN N., WHITEHOUSE M., BAKER G., 2003 - Conserving and promoting parasitoids of *Helicoverpa* in cotton. - Pesticide outlook, 14 (5): 219-221.
- LIMONTA L., DIOLI P., DENTI A., 2003 - Hemiptera present in different plant mixture. - Boll. Zool. agr. Bachic., Ser. II, 35: 55-66.
- LO VERDE G., MASSA B., CALECA V., 1997 - Siepi, bordure e margini di vegetazione spontanea negli agroecosistemi: effetti sulla diversità delle comunità di artropodi. - Naturalista sicil., S. IV, XXI (suppl.): 123-157.
- MACELJSKI M., BALARIN I., 1977 - Beitrag zur Kenntnis natürlicher Feinde der Platanen-Netzwanze (*Corythuca ciliata* (Say), Tingidae, Heteroptera). - Anz. Schädlinke, Pflanzenschutz, Umweltschutz, 50: 135-138.
- MANOJLOVIC B., ZABEL A., KOSTIC M., STANKOVIC S., 2000 - Effect of nutrition of parasites with nectar of melliferous plants on parasitism of the elm bark beetles (Col., Scolytidae). - J. Appl. Entomol., 124: 155-161.

- MANOJLOVIC B., ZABEL A., STANKOVIC S., KOSTIC M., 2001 - Additional diet of the parasitoids (Hymenoptera: Braconidae) and the parasitizing of the Elm Bark Beetle (Coleoptera: Scolytidae). - Anz. Schadlingskunde, 74 (3): 66-71.
- PARENZAN P., PORCELLI F., SINACORI A., 1994 - *Nysius graminicola* (Klt. 1858) e *Oxycarenus hyalinipennis* (Costa 1838), (Rhynchota, Lygaeidae) su kenaf in Basilicata. - Informatore fitopatologico, 44 (1): 62-64.
- PATT J.M., HAMILTON G.C., LASHOMB J.H., 1997 - Foraging success of parasitoid wasps on flowers: interplay of insect morphology, floral architecture and searching behaviour. - Ent. Exp. Appl., 83: 21-30.
- REMUND U., GUT D., BOLLER E.F., 1992 - Beziehungen zwischen Begleitflora und Arthropodenfauna in ostschweizer Rebbergen. Einfluss der botanischen Vielfalt auf ökologische Gleichgewichte. - Schweiz. Z. Obst- u. Wein., 128 (20): 528-540.
- SOLYMOSI P., 1996 - Gyomszabalyozasra hasznalhato donornovenyek. - Novenyvedelem, 32 (1): 23-34 (summary in english).
- SOUTHWOOD T.R.E., LESTON D., 1959 - Land and water bugs of the British Isles. Frederick Warne & Co. LTD, London & New York, pp. 1-544.
- SUMMERS C.G., 1975 - Population fluctuations of selected arthropods in alfalfa: influence of two harvesting practices. - Env. Ent., 5 (1): 103-110.
- TAKESHI T., NUMATA H., 2000 - Effects of feeding on reproduction and overwintering in female adults of *Ooencyrtus nezarae* Ishii (Hymenoptera: Encyrtidae). - Appl. Entomol. Zool., 35(3): 361-367.
- TAMANINI L., 1988 - Tabelle per la determinazione dei più comuni Eterotteri Italiani. - Mem. Soc. Entom. It., 62 (2): 359-471.
- VAN DEEN BOSCH R., LAGACE C.F., STERN V.M., 1967 - The interrelationship of the aphid *Acyrtosiphon pisum*, and its parasite, *Aphidius smithi*, in a stable environment. - Ecology, 48 (6): 993-1000.
- VAN DEEN BOSCH R., MESSNEGER P.S., GUTIERREZ A.P., 1982 - An introduction to Biological Control. New York: Plenum Press, 247 pp.
- VIGNA TAGLIANTI A., AUDISIO P.A., BIONDI M., BOLOGNA M.A., CARPANETO G.M., DE BIASI A., FATTORINI S., PIATTELLA E., SINDACO R., VENCHI A., ZAPPAROLI M., 1999 - A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic region. - Biogeographia, 20: 31-59.
- WACKERS F.L., BJORNSEN A., DORN S., 1996 - A comparison of flowering herbs with respect to their nectar accessibility for the parasitoid *Pimpla turionellae*. Proc. Exper. & Appl. Entomol., N. E. V., Amsterdam, 7: 177-182.
- WAGNER E., 1975 - Die Miridae des Mittelmeerraumes und der Makaronesischen Inseln (Hemiptera, Heteroptera), Teil 3. Ent. Abh. 40. Akad.-Verlagsges. Geest & Portig KG, Leipzig, Germany.
- WARD R.H., PIENKOWSKI R.L., KOK L.T., 1974 - Host specificity of the first-instar of *Ceuthorrhynchidius horridus*, a weevil for biological control of thistles. - J. econ. Entom., 67 (6): 735-737.

PROF. LIDIA LIMONTA, DR NICOLA BONOMELLI - Istituto di Entomologia agraria, Università degli Studi di Milano, Via Celoria 2, I- 0133 Milano. E-mail: lidia.limonta@unimi.it
DR PARIDE DIOLI - Via Valeriana 19 - 23100 Sondrio.

Accepted 4 October 2004