

V. VACANTE, J. GUTIERREZ

Recent advances in agricultural acarology ^(*)

Abstract - The authors describe the recent advances in agricultural acarology and briefly examine the principal research lines that have been involving during the last years various researchers. In particular, the use of informatic application and mathematical models in crop protection against phytophagous mites, the population dynamics, the behaviour of preys and predators, the population genetics and also the molecular biology are discussed.

Riassunto - *Recenti progressi in acarologia agraria.*

Gli autori illustrano le recenti acquisizioni in acarologia agraria ed esaminano brevemente i principali filoni di ricerca che vedono impegnati negli ultimi anni gli studiosi del settore. Si tratta in particolare dell'applicazione dell'informatica e dei modelli matematici nella protezione delle colture dagli acari fitofagi, dell'analisi della dinamica delle popolazioni, del comportamento dei fitofagi e dei predatori, della genetica delle popolazioni, nonché della biologia molecolare.

Key words: advances, agricultural acarology.

INTRODUCTION

As in other branches of biology, fundamental research in modern agricultural acarology has benefited by technical progress in calculation with the development of computing and in biochemistry with that of molecular methods. From this point of view it can be considered that three main lines have been followed. The first consists of the mathematical models developed thanks to progress in data processing. They have led either to the establishment of models applied to crop protection or models for research on population dynamics. The second concerns the analysis of the different aspects of the behaviour of pests and predators. This has developed in particular thanks to progress in data recording methods. The third concerns population genetics. This has expanded considerably with the recent progress in biochemistry and mole-

(*) Paper presented at the XXI International Congress of Entomology in Firenze, August 25-31, 1996.

cular biology. Mites form excellent material for research on population dynamics and the development of models, and many of these appear in recent literature. They are not easy to describe, but we have tried to separate them into crop protection models and research models.

CROP PROTECTION MODELS

Crop protection mathematical models are either tactical or strategic. Tactical models aim at forecasting the scale of attacks, determining the best intervention periods and the quantity of predators to be released or at simulating interactions between prey and predators studied by a stochastic model and pilot experiments (Pearl et al., 1989; Hayes et al., 1993; Sadof & Alexander, 1993). Strategic models approach the population dynamics of phytophagous pests according to the development of the plants attacked (Baumgärtner & Zahner, 1984; Zeng et al., 1985; Wilson et al., 1992). Other authors have examined a simulation model which included the effects of temperature, humidity and predation and coupled to a comprehensive plant-microenvironment model (Berry et al., 1991).

RESEARCH MODELS

Research models are more analytical and explanatory. They examine population dynamics taking into account the plant-mite system, the predator-prey system or the tritrophic plant-mite-predator system.

The first point has been the focus of several studies by Gutierrez et al. (1988), one of the leading initiators of this type of model.

The predator-prey system has been studied according to various special factors, among these the microclimate, with temperature variations or seasonal changes (Rabbinge, 1976; Woolhouse & Harmsen, 1989; Collings et al., 1990), the spatial distribution (Walde et al., 1992) or the dispersal (Sabelis & Laane, 1986; Diekmann et al., 1988; Metz et al., 1988; Janssen & Sabelis, 1992).

As regards the tritrophic system the research has been carried out by different authors (Gutierrez et al., l.c.; Baumgärtner et al., 1988; Wermelinger et al., 1992).

BEHAVIOUR OF PESTS AND PREDATORS

The behaviour of pests and predators involve different aspects; among these the most important are the mating, the spinning, the defence subsocial life, the foraging dispersal, the choice of prey olfactometry and the acaricide resistant populations.

The mating and spinning behaviour of pests and predators have been investigated by various acarologists (Everson & Addicot, 1982; Aponte & McMurtry, 1992; Kumari & Sadana, 1992; Enders, 1993).

Saito (1986), Saito (1990) has studied the sociobiological relations between pests and Yamamura (1987) the biparental defence in a subsocial spider mite.

Foraging dispersal has been examined by Berry et al. (1990) and Zhang et al. (1992). The choice of prey, the predatory and searching behaviour have been studied by Dicke et al. (1986), Dicke (1988), Clements & Harmsen (1990), Santos (1991), Ferragut et al. (1992), Clements & Harmsen (1993) and Sabelis & Van der Weel (1993).

Finally, the population behaviour of acaricide-resistant pests has been investigated by Berry et al. (l.c.), Kolmes et al. (1990) and Kolmes et al. (1994).

POPULATION GENETICS

Much research on isoenzyme variability has been carried out on the Tetranychidae. Initiated by Ward et al. (1982) and by Grafton-Cardwell et al. (1988) in the USA, it was considerably developed in Japan, China and Korea. Investigations have concerned inter and intraspecific polymorphism in Tetranychid and Tenuipalpid mites (Lee, 1988; Kuang & Cheng, 1990; Gotoh & Ishikawa, 1992; Osakabe, 1991a; Osakabe, 1991b; Osakabe 1993; Osakabe et al., 1993; Osakabe & Sakagami, 1993; Hinomoto & Takafuji, 1994; Hinomoto & Takafuji, 1995; Osakabe & Komazaki, 1996; Tsagkarakou et al., 1996b), relationships between Eriophyid mite families (Kuang et al., 1992), polymorphism linked to the diapause status of populations (Gotoh et al., 1991; Osakabe, l.c.) and to acaricidal resistance (Kim & Lee, 1990; Mowry et al., 1996; Tsagkarakou et al., 1996a).

MOLECULAR BIOLOGY

Molecular biology should enable accurate identification of species, the establishment of phylogenies, the discovery of markers for phytophagous and predators and lead to interesting genetic transformation of predators. For the Tetranychid mites are well known the contributions of Kaliszewski et al. (1992), Navajas et al. (1992), Fournier et al. (1994), Navajas et al. (1994), Navajas et al. (1996a) and Navajas et al. (1996b). The Eriophyid identification has been investigated by Fenton et al. (1995). The stable genetic transformations of predators has been treated by Presnail & Hoy (1992). Jeyapragash & Hoy (1995) refer on the complete sequence of a mariner transposable element from a Phytoseiid isolated by an inverse PCR approach.

CONCLUSION

We can generally affirm that the today's agricultural acarology foster great interests not only for the morphological or traditional systematics and for the qualitative settlement of the agroecosystems, but also for the quantitative analysis of the problem, with particular interest for the relation pest-predator-plant and environment. This needs

a better knowledges of biology (*sensu lato*) and involves different competences. From this point of view in the recent years the behaviour of pests and predators, the population dynamics, the crop protection models, the genetic populations and the molecular biology have polarized the interest of many scientists. Mathematical and statistical analysis, biochemical and physiology supports have consented flattering aims.

In conclusion research in agricultural acarology has tended to be less abundant in recent years, but the problems investigated are still there or have been replaced by others. In particular, we are confronted by the problem of resistance to acaricides and modification of mite fauna by the accidental introduction of pests or the intentional introduction of predators. The use of recent research techniques is resulting in rapid progress in the discipline and also highlights the advantages of this biological material for more general research.

REFERENCES

- APONTE O., MCMURTRY J., 1992 - Mating behaviour and reproductive mechanisms of *Amblyseius colimensis* Aponte and McMurtry (Acari: Phytoseiidae). - Bol. Entomol. Venezolana, 7 (1): 1-12.
- BAUMGÄRTNER J., ZAHNER P., 1984 - Simulation experiments with stochastic population models as a tool to explore pest management strategies in an apple-tree mite system (*Panonychus ulmi* Koch, *Tetranychus urticae* Koch). In: CAVALLORO R. (Ed.), Statistical and mathematical methods in population dynamics and pest control, Proc. Meeting E.C. experts' group, Parma, 26-28 oct. 1983, Balkema, Rotterdam: 166-178.
- BAUMGÄRTNER J., GUTIERREZ A.P., KLAY A., 1988 - Elements for Modelling the Dynamics of Tritrophic Population Interactions. - Exp. appl. Acarol. 5 (3-4): 243-263.
- BERRY J.S., HOLTZER T.O., BALL H.J., 1990 - Ambulatory Dispersal of Spider Mites (Acari: Tetranychidae) on Whole, Untreated Maize Plants After Exposure to Fenvalerate and Permethrin. - J. econ. Entomol. 83 (1): 217-220.
- BERRY J.S., HOLTZER T.O., NORMAN J.M., 1991 - MiteSim-a simulation model of the Banks grass mite (Acari: Tetranychidae) and the predatory mite, *Neoseiulus fallacis* (Acari: Phytoseiidae) on maize: model development and validation. - Ecol. Modelling, 53 (3-4): 291-317.
- CLEMENTS D.R., HARMSSEN R., 1990 - Predatory behavior and prey-stage preferences of stigmatid and phytoseiid mites and their potential compatibility in biological control. - Can. Entomol. 122 (3-4): 321-328.
- CLEMENTS D.R., HARMSSEN R., 1993 - Prey preferences of adult and immature *Zetzellia mali* Ewing (Acari: Stigmatidae) and *Typhlodromus caudiglans* Schuster (Acari: Phytoseiidae). - Can. Entomol., 125 (5): 967-969.
- COLLINGS J.B., WOLLKIND D.J., MOODY M.E., 1990 - Outbreaks and oscillations in a temperature-dependent model for a mite predator-prey interaction. - Theor. Popul. Biol., 38 (2): 159-191.
- DICKE M., 1988 - Infochemicals in tritrophic interactions. Origin and function in a system consisting of predatory mites, phytophagous mites and their host plants. - Wageningen, Netherlands, Depart. Entomol. 235 pp.

- DICKE M., SABELIS M.W., GROENEVELD A., 1986 - Vitamin A deficiency modifies response of predatory mite *Amblyseius potentillae* to volatile kairomone of two-spotted spider mite, *Tetranychus urticae*. - J. Chem. Ecol. 12 (6): 1389-1396.
- DIEKMANN O., METZ J.A., SABELIS M.W., 1988 - Mathematical Models of Predator/Prey Plant Interactions in a Patch Environment. - Exp. appl. Acarol. 5 (3-4): 319-342.
- ENDERS M.M., 1993 - The effect of male size and operational sex ratio on male mating success in the common spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae). Anim. Behav. 46(5): 835-846.
- EVERSON P.R., ADDICOTT J.F., 1982 - Mate selection strategies by male mites in the absence of intersexual selection by females: a test of six hypotheses. - Can. J. Zool. 60 (11): 2729-2736.
- FENTON B., MALLOCH G., JONES A.T., AMRINE J.W.JR., GORDON S.C., A'HARA S., MCGAVIN W.J., BIRCH A.N.E., 1995 - Species identification of *Cecidophyopsis* mites (Acari: Eriophyidae) from different *Ribes* species and countries using molecular genetics. - Mol. Ecol. 4(3): 383-387.
- FERRAGUT F., LABORDA R., COSTA-COMELLES J., GARCIA-MARI F., 1992 - Feeding behavior of *Euseius stipulatus* and *Typhlodromus phialatus* on the citrus red mite *Panonychus citri* (Acari: Phytoseiidae, Tetranychidae) - Entomophaga 37 (4): 537-543.
- FOURNIER D., BRIDE J.M., NAVAJAS M., 1994 - Mitochondrial DNA from a spider mite: isolation, restriction map and partial sequence of the cytochrome oxidase subunit I gene. - Genetica 94(1): 73-75.
- GOTOH T., ISHIKAWA Y., TAKAYAMA K., SUZUKI M., 1991 - Differences in esterase zymograms between diapausing and nondiapausing individuals of the hawthorn spider mite, *Tetranychus viennensis* Zacher (Acari: Tetranychidae). - Appl. Entomol. Zool. 26 (1): 153-155.
- GOTOH T., ISHIKAWA Y., 1992 - Separation of *Panonychus ulmi*, *P. thelytokus* and *P. bambuscola* (Acari: Tetranychidae) by esterase zymograms. - Appl. Entomol. Zool. 27 (4): 598-601.
- GRAFTON-CARDWELL E.E., EASH J.A., GRANETT J., 1988 - Isozyme Differentiation of *Tetranychus pacificus* from *T. urticae* and *T. turkestanii* (Acari: Tetranychidae) in Laboratory and Field Populations. - J. econ. Entomol. 81 (3): 770-775.
- GUTIERREZ A.P., WERMELINGER B., SCHULTHESS F., BAUMGÄRTNER J.U., HERREN H.R., ELLIS C.K., YANINEK J.S., 1988 - Analysis of biological control of cassava pests in Africa. I simulation of carbon, nitrogen and water dynamics in cassava. - J. appl. Ecol. 25 (3): 901-920.
- HAYES A.J., WALKER J.T.S., SHAW P.W., WHITE V., 1993 - A decision model for miticide use in apple orchards. - In Proc. Forty Sixth new Zealand plant protection Conference, Christchurch, N.Z. : 162-165.
- HINOMOTO N., TAKAFUJI A., 1994 - Studies on the population structures of the two-spotted mite, *Tetranychus urticae* Koch, by allozyme variability analysis. - Appl. Entomol. Zool. 29 (2): 259-266.
- HINOMOTO N., TAKAFUJI A., 1995 - Genetic changes in the population structure of the two spotted spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae), on vinyl-house strawberries. - Appl. Entomol. Zool. 30(4): 521-528.
- JANSSEN A., SABELIS M.W., 1992 - Phytoseiid life-histories, local predatory-prey dynamics, and strategies for control of tetranychid mites. - Exp. appl. Acarol. 14 (3-4): 233-250.

- JEYAPRAKASH A., HOY M.A., 1995 - Complete sequence of a mariner transposable element from the predatory mite *Metaseiulus occidentalis* isolated by an inverse PCR approach. - *Insect Mol. Biol.* 4 (1): 31-39.
- KALISZEWSKI M.J., TOBOLEWSKI J., SEYOUM S., CHOJNACKI I., KALISZEWSKA M.M., STANTON D.J., COLWELL R.K., 1992 - The polymerase chain reaction and sequencing of mite DNA. - *Int. J. Acarol.* 18 (3): 231-239.
- KIM S.S., LEE S.C., 1990 - Development of acaricidal resistance and esterase isozyme of *Tetranychus urticae* (Acarina: Tetranychidae). - *Korean J. appl. Entomol.* 29 (3): 170-175.
- KOLMES S.A., DENNEHY T.J., FRISICANO L., 1990 - Behavioral aspects of dicofol resistance in the two spotted mite, *Tetranychus urticae*. - *Entomol. Exp. appl.* 57 (3): 259-269.
- KOLMES S.A., DENNEHY T.J., SAM Y., 1994 - Contrasting Behavior of Two-Spotted Spider Mite (Acari: Tetranychidae) on Discontinuous Residues of a Pyrethroid and a Chlorinated Hydrocarbon Acaricide. - *J. econ. Entomol.* 87 (3): 559-565.
- KUANG H.Y., CHENG L.S., 1990 - Studies on the differentiation of two sibling species, *Tetranychus cinnabarinus* (Boisduval) and *T. urticae* Koch. - *Acta entom. Sinica*, 33 (1): 109-116.
- KUANG H.Y., XIE M.C., GONG G.J., 1992 - A preliminary study on the relationship between the families of Eriophyoidea. - *Acta Zootaxon. Sinica*, 17 (3): 294-298.
- KUMARI M., SADANA G.L., 1992 - Mating behaviour of predatory mite, *Amblyseius alstoniae* Gupta (Acari: Phytoseiidae). - *J. Insect Sci.*, 5 (2): 165-166.
- LEE J.S., 1988 - A study on the taxonomy and isozymes of false spider mites (Acarina: Tenuipalpidae) from Korea. *Korean J. Zool.* 31: 147-155.
- METZ J.A.J., SABELIS M.W., KUCHLEIN J.H., 1988 - Sources of Variation in Predation Rates at High Prey Densities: an Analytic Model and a Mite Example. - *Exp. appl. Acarol.* 5 (3-4): 187-205.
- MOWRY T.M., WHALON M.E., CROFT B.A., 1996 - Identification of resistant strains of the predatory mite, *Amblyseius fallacis*, by Polyacrylamide Gel Electrophoresis. (pers. Comm.)
- NAVAJAS M., COTTON D., KREITER S., GUTIERREZ J., 1992 - Molecular approach in spider mites (Acari: Tetranychidae): preliminary data on ribosomal DNA sequences. - *Exp. appl. Acarol.* 15: 211-218.
- NAVAJAS M., GUTIERREZ J., BONATO O., BOLLAND H.R., MAPANGOU DIVASSA S., 1994 - Intra-specific diversity of the cassava green mite *Mononychellus progresivus* (Acari: Tetranychidae) using comparisons of mitochondrial and nuclear ribosomal DNA sequences and cross-breeding. - *Exp. appl. Acarol.* 18: 351-360.
- NAVAJAS M., FOURNIER D., LAGNEL J., GUTIERREZ J., BOURSOT P., 1996a - Mitochondrial COI sequences in mites: evidence for variations in base composition. - *Insect Mol. Biol.* 5 (3): 281-285.
- NAVAJAS M., GUTIERREZ J., LAGNEL J., 1996b - Mitochondrial cytochrome oxidase I in tetranychid mites: a comparison between molecular phylogeny and changes of morphological and life history traits. - *Bull. entom. Res.* 86: 407-417.
- OSAKABE M., 1991a - Genetic analysis of esterases in the Citrus red mite, *Panonychus citri* (McGregor) (Acari: Tetranychidae). - *Appl. Entomol. Zool.* 26 (3): 307-312.
- OSAKABE M., 1991b - Esterase isozyme patterns corresponding to differences in diapause attribute among the populations of the Citrus red mite, *Panonychus citri* (McGregor) (Acari: Tetranychidae). - *Bull. Fruit Tree Res. Sta.* 21: 85-94.
- OSAKABE M., 1993 - Divergence of the northern and south-western populations of *Panonychus mori* Yokohama (Acari: Tetranychidae) in Japan in host range and reproductive compatibility. - *Appl. Entomol. Zool.* 28 (2): 85-94.

- OSAKABE M., SAITO Y., SAKAGAMI Y., 1993 - Protein differences detected by two-dimensional electrophoresis in the species complex of *Schizotetranychus celarius* (Banks). - Exp. appl. Acarol. 17(10): 757-764.
- OSAKABE M., SAKAGAMI Y., 1993 - Estimation of genetic variation in Japanese populations of the Citrus red mite *Panonychus citri* (McGregor) (Acari: Tetranychidae) on the basis of esterase allele frequencies. - Exp. appl. Acarol. 17(10): 749-755.
- OSAKABE M., KOMAZAKI S., 1996 - Differences in esterase isozymes between *Panonychus citri* populations infesting citrus and *Osmanthus*. - Appl. Entomol. Acarol. 20: 113-119.
- PEARL D.K., BARTOSZYNSKI R., HORN D.J., 1989 - A stochastic model for simulation of interactions between phytophagous spider mite and their phytoseiid predators. - Exp. appl. Acarol. 7(2): 143-151.
- PRESNAIL J.K., HOY M.A., 1992 - Stable genetic transformation of a beneficial arthropod, *Metaseiulus occidentalis* (Acari: Phytoseiidae), by a microinjection technique. - Proc. National Academy Sciences United States America 89(16): 7732-7736.
- RABBINGE R., 1976 - Biological control of fruit-tree red spider mite. - Centre Agr. Pub. Doc., Wageningen, 228 pp.
- SABELIS M.W., LAANE W.E.M., 1986 - Regional dynamics of spider mite populations that become extinct locally because of food-source depletion and predation by phytoseiid mites. In: METZ J.A.J., DIEKMANN O. (Eds.), The dynamics of physiologically structure populations. - Lecture notes in Biomathematics, 68. Springer, Berlin: 345-375.
- SABELIS M.W., WEEL J.J. VAN DER, 1993 - Anemotactic responses of the predatory mites, *Phytoseiulus persimilis* Athias-Henriot, and their role in prey finding. - Exp. appl. Acarol. 17: 521-529.
- SADOF C.S., ALEXANDER C.M., 1993 - Limitations of Cost-Benefit-Based Aesthetic Injury Levels for Managing Twospotted Spider Mites (Acari: Tetranychidae). - J. econ. Entomol. 86 (5): 1516-1521.
- SAITO Y., 1986 - Prey kills predator: counter-attack success of a spider mite against its specific phytoseiid predator. - Exp. appl. Acarol. 2: 47-62.
- SAITO Y., 1990 - "Harem" and "non harem" type mating system in two species of subsocial spider mite (Acari, Tetranychidae). - Res. Pop. Ecol. 32 (2): 263-278.
- SANTOS M.A., 1991 - Searching behavior and associational response of *Zetzellia mali* (Acarina: Stigmaeidae). - Exp. appl. Acarol. 11: 81-87.
- TSAGKARAKOU A., NAVAJAS M., LAGNEL J., GUTIERREZ J., PASTEUR N., 1996a - Genetic variability in *Tetranychus urticae* (Acari: Tetranychidae) from Greece: insecticide resistance and isozymes. - J. econ. Entomol. 89 (in press).
- TSAGKARAKOU A., NAVAJAS M., LAGNEL J., PASTEUR N., 1996b - Population structure in the spider mite *Tetranychus urticae* (Acari: Tetranychidae) from Crete based on multiple allozymes. - Heredity 77 (in press).
- WALDE S.J., NYROP J.P., HARDMAN J.M., 1992 - Dynamics of *Panonychus ulmi* and *Typhlodromus pyri*: factors contributing to persistence. - Exp. appl. Acarol. 14: 261-291.
- WARD P.S., BOUSSY I.A., SWINCER D.E., 1982 - Electrophoretic Detection of Enzyme Polymorphism and differentiation in Three Species of Spider Mites (*Tetranychus*) (Acari: Tetranychidae). - Ann. entom. Soc. Am. 75 (6): 595-598.
- WERMELINGER B., CANDOLFI M.P., BAUMGÄRTNER J., 1992 - A model of the European red mite (Acari, Tetranychidae) population dynamics and its linkage to grapevine growth and development. - J. appl. Entomol. 114 (2): 155-166.

- WILSON L.T., CORBETT A., TRICHILO P.J., KERBY T.A., PLANT R.E., GOODELL P.B., 1992 - Strategic and tactical modelling: cotton-spider mite agroecosystem management. - Exp. appl. Acarol. 14: 357-370.
- WOOLHOUSE M.E.J., HARMSSEN R., 1989 - A transition matrix model of European red mite (*Panonychus ulmi*) population dynamics in a managed apple orchards. - Ecol. Model. 46 (3-4): 269-282.
- YAMAMURA N., 1987 - Biparental defence in a subsocial spider mite. - Trends Ecol. Evol. 2 (9): 261-262.
- ZENG Z., LI L.S., ZHOU X.Y., 1985 - Research on the population system and optimal management of the citrus yellow mite (*Eotetranychus kankitus* E.). I. Principal component analysis of the population system of the citrus yellow mite. II. Establishing the sub-models of the population system of the citrus yellow mite. III. Modelling the population system of the citrus yellow mite. IV. Optimal management of the population system of the citrus yellow mite. - J. Southwest Agric. College, China, N°3: 98-123.
- ZHANG Z.Q., SANDERSON J.P., NYROP J.P., 1992 - Foraging time and spatial patterns of predation in experimental population. A comparative study of three mite predator-prey systems (Acari: Phytoseiidae, Tetranychidae). - Oecologia 90 (2): 185-196.

PROF. VINCENZO VACANTE - Dipartimento di Protezione delle Piante. Università della Tuscia,
Via S. Camillo de Lellis, I-01100 Viterbo, Italia

DR. JEAN GUTIERREZ - Laboratoire de Zoologie, INRA - ENSAM - ORSTOM, 2, Place P. Viala,
34060 Montpellier, France.

Accepted 15 May 1997.